

730 Vassar Avenue, Lakewood, NJ 08701 800.445.9603 \* 732.942.1660 fax 800.368.3108 \* 732.942.9270 www.worthington-biochem.com

**Date:** 10/6/20

## **Certificate of Analysis**

Product Trypsin Inhbitor, Ovomucoid

Source Egg White

Country of

Manufacture USA

All products from animal sources are produced from starting material of USDA-approved origin, collected in United States Department of Agriculture (USDA) or equivalent approved facilities, inspected to be free of disease

and suitable for exportation. Certificates of Origin are available upon request.

Storage Store at 2-8°C.

Code OI

Lot Number

Re-Assay Date

Description Mucoprotein and antitryptic factor of egg white described by Lineweaver and Murray, J. Biol. Chem.,

171, 565 (1947). A dialyzed, dried powder.

Unit Definition One milligram inhibits at least 1.2 milligrams of trypsin, Code TRL, with at least 180 Units per

milligram. (One Unit TRL hydrolyzes one micromole of TAME per minute at 25°C, pH 8.2, in the

presence of 0.1M calcium.)

<u>Parameter</u>	Result	Acceptance Criteria
mg TRL inhibited by 1 mg	1.37	1 mg inhibits at least 1.20 mg trypsin,
		Code: TRL
% Protein	64%	Report assay value.
Endotoxin EU/mg dw	< 0.5	Report Assay Value.