



730 Vassar Avenue, Lakewood, NJ 08701  
800.445.9603 \* 732.942.1660 fax 800.368.3108 \* 732.942.9270  
[www.worthington-biochem.com](http://www.worthington-biochem.com)

### Certificate of Analysis

Product Papain, Suspension  
Source Carica papaya Latex  
Country of Manufacture USA  
Storage Store at 2-8°C.  
Code PAP  
Lot Number  
Re-Assay Date  
Description Supplied as a two times crystalline suspension in 50mM sodium acetate, pH 4.5. To ensure full activity, the enzyme should be incubated in a solution containing 1.1mM EDTA, 0.067mM mercaptoethanol and 5.5mM cysteine-HCl for 30 minutes. It is recommended that the enzyme be 0.22 micron filtered after dissolution and prior to use.  
Unit Definition One Unit hydrolyzes one micromole of benzoyl-L-arginine ethyl ester per minute at 25°C, pH 6.2, after activation in a solution containing 1.1mM EDTA, 0.067mM mercaptoethanol and 5.5mM cysteine-HCl for 30 minutes.

<u>Parameter</u>	<u>Result</u>	<u>Acceptance Criteria</u>
u/mgP	30.1	Activates to at least 20.0 units per mg protein
mgP/ml	41.9	Report assay value.
u/ml	1,260	Report assay value.
Endotoxin EU/ml	1.15	Report Assay Value.

Approved by: Thomas R. Ryan, Quality Control Manager (signature on file)

Date: 10/15/20